

(19) World Intellectual Property Organization  
International Bureau







(43) International Publication Date  
7 June 2001 (07.06.2001)

PCT

(10) International Publication Number  
**WO 01/40180 A2**

- (51) International Patent Classification<sup>7</sup>: C07D 207/16, A61K 31/401
- (74) Agent: GEERING, Keith, Edwin; Reddie & Grose, 16 Theobalds Road, London WC1X 8PL (GB).
- (21) International Application Number: PCT/GB00/04572
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date:  
30 November 2000 (30.11.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (30) Priority Data:  
9928330.1 30 November 1999 (30.11.1999) GB
- (71) Applicant (*for all designated States except US*): FER-RING BV [NL/NL]; Polaris Avenue 144, NL-2132 JX Hoofddorp (NL).
- (72) Inventor; and
- (75) Inventor/Applicant (*for US only*): EVANS, David, Michael [GB/GB]; 114 Adelaide Road, St. Denys, Southampton SO17 2HX (GB).
- Published:**  
— Without international search report and to be republished upon receipt of that report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: NOVEL ANTIDIABETIC AGENTS



(57) Abstract: Compounds which are 1-(2'-aminoacyl)-2-cyanopyrrolidine derivatives according to general formula (1) are DP-IV inhibitors for treatment of impaired glucose tolerance or type 2 diabetes; wherein A is selected from groups (2, 3 and 4); X is selected from aminoacyl groups corresponding to the natural amino acids, acyl groups R<sup>3</sup>CO, groups R<sup>4</sup>COOC(R<sup>5</sup>)(R<sup>6</sup>)OCO, methoxycarbonyl, ethoxycarbonyl and benzyloxycarbonyl; R<sup>1</sup> is selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl residues, (CH<sub>2</sub>)<sub>a</sub>NHW<sup>1</sup>, (CH<sub>2</sub>)<sub>b</sub>COW<sup>2</sup>, (CH<sub>2</sub>)<sub>c</sub>OW<sup>3</sup>, CH(Me)OW<sup>4</sup>, (CH<sub>2</sub>)<sub>d</sub>-C<sub>6</sub>H<sub>4</sub>-W<sup>5</sup> and (CH<sub>2</sub>)<sub>e</sub>SW<sup>6</sup>, where a is 2-5, b is 1-4, c is 1-2, d is 1-2, e is 1-3, W<sup>1</sup> is COW<sup>6</sup>, CO<sub>2</sub>W<sup>6</sup> or SO<sub>2</sub>W<sup>6</sup>, W<sup>2</sup> is OH, NH<sub>2</sub>, OW<sup>6</sup> or NHW<sup>6</sup>, W<sup>3</sup> is H or W<sup>6</sup>, W<sup>4</sup> is H or W<sup>6</sup>, W<sup>5</sup> is H, OH or OMe, and W<sup>6</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, optionally substituted phenyl, optionally substituted heteroaryl or benzyl and R<sup>2</sup> is selected from H and (CH<sub>2</sub>)<sub>n</sub>-C<sub>5</sub>H<sub>3</sub>N-Y, where n is 2-4 and Y is H, F, Cl, NO<sub>2</sub> or CN, or R<sup>1</sup> and R<sup>2</sup> together are -(CH<sub>2</sub>)<sub>p</sub>-where p is 3 or 4; R<sup>3</sup> is selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl and phenyl; R<sup>4</sup> is selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl and optionally substituted phenyl; R<sup>5</sup> and R<sup>6</sup> are each independently selected from H and C<sub>1</sub>-C<sub>6</sub> alkyl or together are -(CH<sub>2</sub>)<sub>m</sub>-, where m is 4-6; R<sup>7</sup> is selected from pyridyl and optionally substituted phenyl; R<sup>8</sup> is selected from H and C<sub>1</sub>-C<sub>3</sub> alkyl; and R<sup>9</sup> is selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy and phenyl.

## Novel Antidiabetic Agents

The present invention relates to a series of novel compounds that are useful for the treatment of type 2 diabetes, impaired glucose tolerance and certain other conditions.


### Background to the Invention

The enzyme dipeptidyl peptidase IV (EC.3.4.14.5, herein abbreviated as DP-IV, and elsewhere also known as DPP-IV or DAP-IV) is thought to be involved in the regulation of the activities of several hormones. One such hormone is glucagon-like peptide 1 (GLP-1), which is involved in the regulation of post-prandial blood glucose levels, and which is converted from its active form, GLP-1(5-36), to the inactive GLP-1(7-36) by DP-IV. In cases of type 2 diabetes and impaired glucose tolerance, where hyperglycaemia can lead to tissue damage, it would be advantageous to potentiate the effect of endogenous GLP-1. Accordingly, inhibitors of DP-IV have been proposed as candidate drugs for the treatment of type 2 diabetes and impaired glucose tolerance. For example, Demuth *et al.* (WO97/40832) disclose the effect of N-isoleucyl-pyrrolidine on blood glucose levels in a relevant animal model. This compound, however, may not be sufficiently potent to be a viable therapeutic agent. More potent inhibitors of DP-IV are disclosed by Jenkins *et al.* (WO95/15309) and by Villhauer (WO98/19998), but they tend to be unstable and to cyclize in solution. This instability would lead to difficulties in preparing and storing material of adequate quality for human therapeutic use. Therefore, there remains a need for an agent that inhibits DP-IV *in vivo*, but which is stable enough for commercial manufacture.

**Brief Description of the Invention**

We have now found a series of derivatives that are chemically stable, but which undergo metabolic activation after administration to a human subject to liberate highly potent inhibitors of DP-IV. In the art, such derivatives are generally termed prodrugs. The compounds of the present invention are useful for the treatment of type 2 diabetes and impaired glucose tolerance, as well as other conditions wherein the potentiation of the action of a hormone normally inactivated by DP-IV gives a therapeutic benefit.

The compounds according to the invention are 1-(2'-aminoacyl)-2-cyanopyrrolidine derivatives according to general formula 1:




wherein A is selected from groups 2, 3 and 4; X is selected from an aminoacyl group corresponding to one of the natural amino acids, an acyl group  $R^3CO$ , a group  $R^4COOC(R^5)(R^6)OCO$ , methoxycarbonyl, ethoxycarbonyl and benzyloxycarbonyl;  $R^1$  is selected from H, a  $C_1 - C_6$  alkyl residue,  $(CH_2)_aNHW^1$ ,  $(CH_2)_bCOW^2$ ,  $(CH_2)_cOW^3$ ,  $CH(Me)OW^4$ ,  $(CH_2)_d-C_6H_4-W^5$  and  $(CH_2)_eSW^6$ , where a is 2-5, b is 1-4, c is 1-2, d is 1-2, e is 1-3,  $W^1$  is  $COW^6$ ,  $CO_2W^6$  or  $SO_2W^6$ ,  $W^2$  is OH,  $NH_2$ ,  $OW^6$  or  $NHW^6$ ,  $W^3$  is H or  $W^6$ ,  $W^4$  is H or  $W^6$ ,  $W^5$  is H, OH or OMe, and  $W^6$  is  $C_1 - C_6$  alkyl, optionally substituted phenyl, optionally substituted heteroaryl or benzyl and  $R^2$  is selected from H and  $(CH_2)_n-C_5H_3N-Y$ , where n is 2-4 and Y is H, F, Cl,  $NO_2$  or CN, or  $R^1$  and  $R^2$  together are  $-(CH_2)_p-$  where p is 3 or 4;  $R^3$  is selected from H,  $C_1 - C_6$  alkyl and phenyl;  $R^4$  is selected from H,  $C_1 - C_6$  alkyl, benzyl and optionally substituted phenyl;  $R^5$  and  $R^6$  are each independently selected from H and  $C_1 - C_6$  alkyl or together are  $-(CH_2)_m-$ , where m is 4-6;  $R^7$  is selected from pyridyl and optionally substituted phenyl;  $R^8$  is selected from H and  $C_1 - C_3$  alkyl; and  $R^9$  is selected from H,  $C_1 - C_6$  alkyl,  $C_1 - C_6$  alkoxy and phenyl.

The present invention relates to the novel compounds as defined above, to pharmaceutical compositions in which at least one active agent is such a compound, to the use of such compositions for the treatment of certain medical conditions, and to a method of treatment in which the compounds of the invention are administered to a subject in need of treatment.


## Detailed Description of the Invention

In a first aspect, the present invention comprises a series of novel compounds that are prodrugs of therapeutically useful inhibitors of DP-IV. The compounds of the present invention are 1-(2'-aminoacyl)-2-cyanopyrrolidine derivatives according to general formula 1 below.




1


In this general formula, A is a group selected from 2, 3 and 4.



2



3



4

The dashed bond (broken line) indicates the covalent bond that links the nitrogen atom of A to 1.

The group X is an acyl or oxycarbonyl group. Suitable groups are:

(i) amino acyl groups corresponding to one of the natural amino acids alanine (Ala), arginine (Arg), asparagine (Asn), aspartic acid (Asp), cysteine (Cys), glutamine (Gln), glutamic acid (Glu), glycine (Gly), histidine (His), isoleucine (Ile), leucine (Leu), lysine (Lys), methionine (Met), phenylalanine (Phe), proline (Pro), serine (Ser), threonine (Thr), tryptophan (Trp), tyrosine (Tyr) and valine (Val);

(ii) acyl groups  $R^3CO$ , where  $R^3$  is a hydrogen atom, a  $C_1 - C_6$  alkyl group or a phenyl group;

(iii) acyloxymethyleneoxycarbonyl groups  $R^4COOC(R^5)(R^6)OCO$ , where  $R^4$  is a hydrogen atom, a  $C_1 - C_6$  alkyl group, a benzyl group, or a phenyl group which may further be substituted with a  $C_1 - C_3$  group, and  $R^5$  and  $R^6$  are each independently a hydrogen atom or a  $C_1 - C_6$  alkyl group or  $R^5$  and  $R^6$  together are a polymethylene unit  $-(CH_2)_m-$ , where  $m$  is an integer of 4-6; and

(iv) methoxycarbonyl, ethoxycarbonyl and benzyloxycarbonyl.

$R^1$  is the side-chain of a naturally occurring amino acid, or an analogue thereof. More specifically,  $R^1$  is selected from a hydrogen atom,  $C_1 - C_6$  alkyl residues,  $(CH_2)_aNHW^1$ ,  $(CH_2)_bCOW^2$ ,  $(CH_2)_cOW^3$ ,  $CH(Me)OW^4$ ,  $(CH_2)_d-C_6H_4-W^5$  and  $(CH_2)_eSW^6$ , where  $a$  is an integer of 2-5,  $b$  is an integer of 1-4,  $c$  is 1 or 2,  $d$  is 1 or 2,  $e$  is an integer of 1-3,  $W^1$  is  $COW^6$ ,  $CO_2W^6$  or  $SO_2W^6$ ,  $W^2$  is  $OH$ ,  $NH_2$ ,  $OW^6$  or  $NHW^6$ ,  $W^3$  is  $H$  or  $W^6$ ,  $W^4$  is  $H$  or  $W^6$ ,  $W^5$  is  $H$ ,  $OH$  or  $OMe$ , and  $W^6$  is  $C_1 - C_6$  alkyl, optionally substituted phenyl, optionally substituted heteroaryl or benzyl. Suitable optional substituents on the heteroaryl and phenyl groups include  $C_1 - C_3$  alkyl and  $C_1 - C_3$  alkoxy groups as well as fluorine and chlorine atoms. Up to two such substituents may be present.

$R^2$  is a hydrogen atom or a group  $-(CH_2)_nNH-C_5H_3N-Y$ , where  $n$  is an integer of 2-4,  $C_5H_3N$  is a divalent pyridyl moiety, and  $Y$  is a hydrogen atom, a halogen atom such as a fluorine or chlorine atom, a nitro group or a cyano group.

Alternatively,  $R^1$  and  $R^2$  together may be  $-(CH_2)_p-$  where  $p$  is 3 or 4.

R<sup>7</sup> is selected from pyridyl and optionally substituted phenyl. Suitable optional substituents include C<sub>1</sub> – C<sub>3</sub> alkyl groups, C<sub>1</sub> – C<sub>3</sub> alkoxy groups, halogen atoms, nitro groups, cyano groups and carboxy groups. Up to two such substituents may be present.

R<sup>8</sup> is a hydrogen atom or a C<sub>1</sub> – C<sub>3</sub> alkyl group.


R<sup>9</sup> is a hydrogen atom, a C<sub>1</sub> – C<sub>6</sub> alkyl group, a C<sub>1</sub> – C<sub>6</sub> alkoxy group or a phenyl group.

In the context of the present disclosure, "alkyl" includes straight-chain and branched alkyl groups as well as cycloalkyl groups. For example, C<sub>1</sub> – C<sub>6</sub> alkyl includes methyl, ethyl, isopropyl, *tert*-butyl, neopentyl and cyclohexyl groups. Also, "heteroaryl" is intended to include monocyclic five- and six-membered aromatic rings that include from one to three heteroatoms selected from nitrogen, oxygen and sulphur. For example, heteroaryl includes pyrrolyl, pyridyl, furyl, thienyl, imidazolyl, thiazolyl, isoxazolyl, thiadiazolyl, pyrimidyl and pyrazinyl.

Certain of the compounds of the present invention have acidic or basic properties and so can exist as salts. Insofar as such salts are non-toxic and otherwise pharmaceutically acceptable, they are included within the scope of the invention. Examples of such salts include, but are not limited to, the acetate, hydrochloride, sulphate, phosphate and benzoate salts of basic compounds, and the sodium, potassium and tetra-alkyl ammonium salts of acidic compounds.

Except when R<sup>1</sup> is H, compounds according to general formula 1 have two stereogenic centres (asymmetric carbon atoms), shown below as C\*. The stereochemistry at these two positions is preferably the one illustrated. Certain embodiments of R<sup>1</sup> and X allow for further stereogenic centres to be introduced,

and so the compounds of the invention may exist as epimers, including diastereomers. All such optical isomers, including mixtures of such optical isomers, are considered to be within the scope of the invention.



In a preferred embodiment, the present invention comprises a compound according to general formula 1 in which  $R^1$  is other than H and  $R^2$ , where present, is H. In a more preferred embodiment,  $R^1$  is  $C_1 - C_6$  alkyl.

In another preferred embodiment, the present invention comprises a compound according to general formula 1 in which  $R^1$  is H and A is selected from groups according to general formulae 2 and 4 with  $R^2$  being  $-(CH_2)_nNH-C_5H_3N-Y$ . In a more preferred embodiment n is 2 and Y is CN. In a most preferred embodiment, the NH group is at the 2-position and the CN group is at the 5-position of the pyridine ring.

In another preferred embodiment, the present invention comprises a compound according to general formula 1 in which A is a group according to general formula 2 and X is an amino acyl group. In one more preferred embodiment, X is an amino acyl group corresponding to a basic amino acid such as lysine or arginine, and most preferably arginine. In another more preferred embodiment, X is an amino acyl group corresponding to glycine.

In another preferred embodiment, the present invention comprises a compound according to general formula 1 in which A is a group according to general formula 2



and X is a group  $R^4\text{COOC}(R^5)(R^6)\text{OCO}$ . In one more preferred embodiment,  $R^4$  is  $C_1 - C_6$  alkyl. In another more preferred embodiment, one of  $R^5$  and  $R^6$  is H and the other is methyl. Most preferably,  $R^4$  and one of  $R^5$  and  $R^6$  are methyl and the other of  $R^5$  and  $R^6$  is H.

In another preferred embodiment, the present invention comprises a compound according to general formula 1 in which A is a group according to general formula 2 and X is a methoxycarbonyl group.

In another preferred embodiment, the present invention comprises a compound according to general formula 1 in which A is a group according to general formula 3.


In another preferred embodiment, the present invention comprises a compound according to general formula 1 in which A is a group according to general formula 4. In a more preferred embodiment  $R^8$  is  $C_1 - C_3$  alkyl, and most preferably it is methyl. In another more preferred embodiment,  $R^9$  is  $C_1 - C_3$  alkyl or  $C_1 - C_3$  alkoxy, and most preferably it is methyl or methoxy.

Compounds that incorporate the features of more than one of these preferred embodiments are particularly preferred. A most preferred embodiment of the present invention is a compound selected from:

- (2S)-1-((2'S)-2'-(1''-acetoxyethoxycarbonylamino)-3',3'-dimethylbutanoyl)-pyrrolidine-2-carbonitrile;
- (2S)-1-(N'-(1''-acetoxyethoxycarbonyl)isoleucyl)pyrrolidine-2-carbonitrile;
- (2S)-1-(N'-(methoxycarbonyl)isoleucyl)pyrrolidine-2-carbonitrile;
- (2S)-1-((N')-(4''-oxopent-2''-en-2''-yl)isoleucyl)pyrrolidine-2-carbonitrile;
- (2S)-1-(glycylisoleucyl)pyrrolidine-2-carbonitrile;
- (2S)-1-(arginylisoleucyl)pyrrolidine-2-carbonitrile;


(2S)-1-((2'S)-2'-(acetoxymethoxycarbonylamino)-3',3'-dimethylbutanoyl)-pyrrolidine-2-carbonitrile;  
 (2S)-1-((2'S)-2'-(1''-acetoxyethoxycarbonylamino)-2'-cyclohexylacetyl)-pyrrolidine-2-carbonitrile;  
 (2S)-1-((2'S)-2'-(1''-acetoxyethoxycarbonylamino)-4',4'-dimethylpentanoyl)-pyrrolidine-2-carbonitrile;  
 (2S)-1-(N'-(1''-acetoxyethoxycarbonyl)-O'-*tert*-butylserinyl)pyrrolidine-2-carbonitrile;  
 (2S)-1-(N<sup>α</sup>-(1'-acetoxyethoxycarbonyl)-N<sup>ω</sup>-*p*-toluenesulphonyllysiny)-pyrrolidine-2-carbonitrile;  
 (2S)-1-(N-(1'-acetoxyethoxycarbonyl)-N-(2''-(5'''-cyanopyridin-2'''-ylamino)-ethyl)glyciny)pyrrolidine-2-carbonitrile;  
 (2S)-1-(N'-(benzyloxycarbonyl)-O'-*tert*-butylthreoninyl)pyrrolidine-2-carbonitrile;  
 (2S)-1-(S'-*tert*-butyl-N'-(ethyloxycarbonyl)cysteinyl)pyrrolidine-2-carbonitrile;  
 (2S)-1-(N<sup>ω</sup>-acetyl-N<sup>α</sup>-benzoyllysiny)pyrrolidine-2-carbonitrile; and  
 (2S)-1-(N<sup>α</sup>-(acetyl)-N<sup>ω</sup>-(benzyloxycarbonyl)ornithiny)pyrrolidine-2-carbonitrile

The compounds according to the present invention can be prepared by standard techniques that are well known in the field of organic chemistry. In many cases, a suitable starting material is an amine according to general formula 5, in which R<sup>1</sup> and R<sup>2</sup> have the same meaning as defined previously.




The synthesis of such compounds is described in, for example, Jenkins *et al.* (WO95/15309), Villhauer (WO98/19998), Ashworth *et al.* (Bioorg. Med. Chem. Lett. 1996, 6(10), 1163-66) and Li *et al.* (Arch. Biochem. Biophys. 1995, 323(1), 148-54).

Compounds not explicitly described in these publications can be made by routine modification of the methods given therein. The steps involved in the preparation of the compounds of the invention from compounds according to general formula 5 depend on the nature of the group A.



Scheme 1



Scheme 1 illustrates the preparation of these compounds in two steps. Chn represents the side chain of an amino acid. Depending on the amino acid being used, Chn may be H (for glycine), CH<sub>3</sub> (alanine), (CH<sub>3</sub>)<sub>2</sub>CH (valine), (CH<sub>3</sub>)<sub>2</sub>CHCH<sub>2</sub>


(leucine),  $\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)$  (isoleucine),  $\text{C}_6\text{H}_5\text{CH}_2$  (phenylalanine),  $\text{HOC}_6\text{H}_4\text{CH}_2$  (tyrosine),  $\text{C}_8\text{H}_6\text{NCH}_2$  (tryptophan),  $\text{HOOCCH}_2$  (aspartic acid),  $\text{HOOCCH}_2\text{CH}_2$  (glutamic acid),  $\text{H}_2\text{NOCCH}_2$  (asparagine),  $\text{H}_2\text{NOCCH}_2\text{CH}_2$  (glutamine),  $\text{HOCH}_2$  (serine),  $\text{CH}_3\text{CH}(\text{OH})$  (threonine),  $\text{HSCH}_2$  (cysteine),  $\text{CH}_3\text{SCH}_2\text{CH}_2$  (methionine),  $\text{C}_3\text{H}_3\text{N}_2\text{CH}_2$  (histidine),  $\text{H}_2\text{N}(\text{CH}_2)_4$  (lysine) and  $\text{H}_2\text{NC}(\text{:NH})(\text{CH}_2)_3$  (arginine). As will be understood by those familiar with the practice of peptide chemistry, several of these side chains contain functional groups that are reactive under the conditions necessary to effect the condensation of the two fragments. These functional groups must be protected with an appropriate masking group. Such groups are described in, for example, "Protective Groups in Organic Synthesis", T.W. Greene, Wiley-Interscience, 1981.  $\text{Chn}^*$  therefor represents the same side chains but with any necessary protecting groups.

Similarly, PG represents a protecting group for an amino function.


The 1-(2'-aminoacyl)-2-cyanopyrrolidine **5** can be condensed with the appropriately protected amino acid **6** to give intermediate **7** using a variety of conditions that are well known in the field of peptide chemistry. Generally, the two components are dissolved in an appropriate solvent, which is normally an aprotic solvent such as dichloromethane or dimethylformamide or a mixture of these, and the solution is cooled to  $0^\circ\text{C}$  or below. One or two equivalents of an amine base such as diisopropylethylamine or dimethylaminopyridine may be added to the solution. A condensing agent is then added and the mixture is stirred until the starting materials have been consumed, as indicated by, for example, analytical thin layer chromatography. If the reaction is slow, it may be advisable to allow the mixture to warm up to ambient temperature to accelerate the process. Suitable condensing agents include DCC (dicyclohexylcarbodiimide), BOP ((benzotriazol-1-yloxy)-tris(dimethylamino)phosphonium hexafluorophosphate), PyBOP® ((benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate), PyBroP® (bromo-tripyrrolidinophosphonium hexafluorophosphate) and HBTU (O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate).

Deprotection of intermediate 7 gives the target compound 1<sup>A</sup> (i.e. a compound of general formula 1 with A according to general formula 2 and X an amino acyl residue).


Suitably protected proline may be used instead of 6 to give the analogous compound with X being a prolyl residue. All the protected amino acids are items of commerce.




Scheme 2




Scheme 2 illustrates the preparation of these compounds. The starting material 5 is treated with an acyl chloride or an anhydride in an aprotic solvent and in the presence of an amine base such as described above, to give the product 1<sup>B</sup>. When R<sup>3</sup> = H it is not possible to use the acyl chloride or anhydride. In this case a mixed anhydride is used. The reagent can conveniently be prepared from formic acid and acetic anhydride.




Scheme 3




Scheme 3 illustrates the preparation of these compounds. The starting material **5** is treated with a *p*-nitrophenyl carbonate **8** in an aprotic solvent and in the presence of an amine base such as described previously to give the product **1<sup>c</sup>**. The carbonate is prepared according to the method described by Alexander *et al.*, J. Med. Chem. 31, 318, 1988.




Scheme 4



Scheme 4 illustrates the preparation of these compounds. The starting material 5 is treated with a chloroformate in an aprotic solvent and in the presence of an amine base such as described previously to give the product 1<sup>D</sup>. Since benzyl chloroformate (BnOC(=O)Cl) is not very stable, it may conveniently be replaced by benzyl 1-succinimidyl carbonate (BnOC(=O)ONSu). This and all the chloroformates are items of commerce.




Scheme 5




Scheme 5 illustrates the preparation of these compounds. The starting material **5A** (i.e. a compound according to general formula **5** with  $R^2 = H$ ) is reacted with an aldehyde **9** in the presence of an acidic catalyst such as, for example, *p*-toluenesulphonic acid. The reaction is performed in a solvent such as cyclohexane or toluene at an elevated temperature such as at the boiling point of the solvent. Water is removed continuously, either by azeotropic distillation or with a desiccating agent such as activated molecular sieves.

The aldehydes **9** are items of commerce.



Scheme 6






Scheme 6 illustrates the preparation of these compounds. The starting material **5** is reacted with an 1,3-dicarbonyl compound **10** in an aprotic solvent and in the presence of an amine base, as previously described, at ambient temperature.

The dicarbonyl compounds **10** are either items of commerce or may be prepared according to well-established procedures.

Other synthetic routes are, of course, possible. In general, they differ from those described above in the order in which steps are performed. Two examples are illustrated in Scheme 7.

Scheme 7



Intermediate **11** is prepared according to the methods previously described. It can be condensed with prolineamide (**12**) according to the methods described in Scheme 1 to give the intermediate **13**. This can be dehydrated by treatment with

trifluoroacetic anhydride to give the target compound. Alternatively, intermediate 11 can be condensed with prolinenitrile (14) to give the target compounds directly.

Compounds according to general formula 1 (the compounds of the present invention) are metabolised in the body to give compounds according to general formula 5. These metabolites are inhibitors of DP-IV.

Scheme 7



As discussed previously, inhibitors of DP-IV are believed to be useful in the treatment of certain medical conditions. Accordingly, the compounds of the present invention are useful in the treatment of these same medical conditions. In particular, the compounds of the present invention are useful in the treatment of impaired glucose tolerance and type 2 diabetes. They can also be useful in the treatment of reproductive disorders such as infertility due to polycystic ovary syndrome. A further use is in the treatment of growth hormone insufficiency leading to small stature. Other medical conditions are also included within the scope of the invention.

For use in the treatment of these disorders, the compounds of the invention will generally be included in a pharmaceutical composition and formulated appropriately for the intended route of administration. Such compositions comprise a second aspect to the present invention. The pharmaceutical composition may include other such pharmaceutically acceptable excipients as are generally known in the art, such as bulking agents, diluents, dispersants, preservatives, colouring and flavouring agents and the like. . The choice of the

excipients will depend on the manner in which the composition is to be formulated and administered. The composition may be administered by the routes generally known in the art. For example, the composition may be formulated as a tablet, capsule, syrup or powder for oral administration, as a lozenge or wafer for sublingual or buccal administration, as a suppository for rectal or vaginal administration, as a solution, suspension or powder for nasal administration, as a cream or lotion for topical administration, as a patch for transdermal administration, or as a solution or suspension for subcutaneous, intramuscular or intravenous injection. Injectable forms may include encapsulated and other controlled-release formulations as are known in the art to be suitable for depot administration. A preferred composition is a tablet for oral administration.

In a third aspect, the invention comprises a method of treatment of glucose intolerance or type 2 diabetes wherein a person in need of such treatment is administered a therapeutically effective amount of a compound as described above. The dosing regimen will generally be decided by the treating physician, taking into account the particular characteristics of the patient. The dose will typically be from 1mg to 500mg once per day or up to four times per day.


The foregoing general description is further illustrated below in a number of examples. These are intended to demonstrate the implementation of the invention, but they do not in any way limit the scope of what has been described hereto.

## EXAMPLES

Solvents and reagents were generally used as supplied without further purification.

The structures of all intermediates were confirmed by <sup>1</sup>H NMR. Final products were further characterised by mass spectroscopy and/or elemental analysis.

Example 1 – (2S)-1-((2'S)-2'-(1''-Acetoxyethoxycarbonylamino)-3',3'-dimethylbutanoyl)pyrrolidine-2-carbonitrile




A solution of (2S)-1-((2'S)-2'-amino-3',3'-dimethylbutanoyl)pyrrolidine-2-carbonitrile hydrochloride (180mg, 0.73mmol; prepared according to Jenkins *et al.*, WO95/15309),  $\alpha$ -acetoxyethyl *p*-nitrophenyl carbonate (220mg, 0.82mmol; prepared according to Alexander *et al.*, J. Med. Chem. 31, 318, 1988) and triethylamine (90mg, 0.90mmol) in dichloromethane (25ml) was stirred at room temperature for 18 hours. After this time the solvent was removed *in vacuo* and the residue was taken up in ethyl acetate (70ml). This solution was washed with 0.3M KHSO<sub>4</sub>, sat. NaHCO<sub>3</sub>, water and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by flash chromatography (eluant EtOAc:Pet. Ether 60-80°C; 3:7) yielding a white solid identified as the title compound (170mg, 0.50mmol, 68%).

MS:- ESI {M+H}<sup>+</sup> = 340.2

<sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  1.02, 1.03 (9H, 2xs), 1.42-1.46 (3H, m), 2.03, 2.05 (3H, 2xs), 2.15-2.25 (4H, m), 3.69-3.76 (2H, m), 4.23-4.28 (1H, m), 4.77-4.79 (1H, m), 5.43 (1H, d, J = 9.5Hz), 6.73-6.77 (1H, m) ppm

Example 2 – (2S)-1-(N'-(1''-Acetoxyethoxycarbonyl)isoleucyl)pyrrolidine-2-carbonitrile




A solution of (2S)-1-(isoleucyl)pyrrolidine-2-carbonitrile hydrochloride (500mg, 2.04mmol; prepared according to Jenkins *et al.*, WO95/15309),  $\alpha$ -acetoxyethyl *p*-nitrophenyl carbonate (610mg, 2.27mmol; prepared according to Alexander *et al.*, *J. Med. Chem.* 31, 318, 1988) and triethylamine (250mg, 2.50mmol) in dichloromethane (40ml) was stirred at room temperature for 18 hours. After this time the solvent was removed *in vacuo* and the residue was taken up in ethyl acetate (70ml). This solution was washed with 0.3M KHSO<sub>4</sub>, sat. NaHCO<sub>3</sub>, water and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by flash chromatography (eluant EtOAc:Pet. Ether 60-80°C; 3:7) yielding a colourless oil identified as the title compound (480mg, 1.42mmol, 70%).

MS:- ESI {M+H}<sup>+</sup> = 340.0

<sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  0.86-0.89 (6H, m), 0.92-0.97 (1H, m), 1.41-1.45 (3H, m), 1.50-1.80 (2H, m), 2.02 (3H, d, J = 5.2Hz), 2.14 -2.27 (4H, m), 3.60-3.75 (2H, m), 4.23-4.26 (1H, t, J = 7.6Hz), 4.77 (1H, d, J = 2.3Hz), 5.30-5.50 (1H, m), 6.73-6.77 (1H, m) ppm

## Example 3 – (2S)-1-(N'-(Methoxycarbonyl)isoleucyl)pyrrolidine-2-carbonitrile




A solution of (2S)-1-(isoleucyl)pyrrolidine-2-carbonitrile hydrochloride (300mg, 1.22mmol; prepared according to Jenkins *et al.*, WO95/15309), methyl chloroformate (125mg, 1.3mmol) and triethylamine (150mg, 1.50mmol) in dichloromethane (40ml) was stirred at room temperature for 18 hours. After this time the solvent was removed *in vacuo* and the residue was taken up in ethyl acetate (70ml). This solution was washed with 0.3M KHSO<sub>4</sub>, sat. NaHCO<sub>3</sub>, water and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by flash chromatography (EtOAc:Pet. Ether 60-80°C; 4:6) yielding a colourless oil identified as the title compound (310mg, 1.16mmol, 95%).

MS:- ESI {M+H}<sup>+</sup> = 268.2

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 0.85-0.95 (6H, m), 1.10-1.25 (1H, m), 1.54-1.77 (2H, m), 2.11-2.26 (4H, m), 3.62 (3H, s), 3.66-3.79 (2H, m), 4.21 (1H, t, J = 9.2Hz), 4.74-4.78 (1H, m), 5.30 (1H, d, J = 9.1Hz) ppm.

**Example 4 – (2S)-1-((N')-(4''-Oxopent-2''-en-2''-yl)isoleucyl)pyrrolidine-2-carbonitrile**




A solution of (2S)-1-(isoleucyl)pyrrolidine-2-carbonitrile hydrochloride (150mg, 0.61mmol; prepared according to Jenkins *et al.*, WO95/15309), 2,4-pentanedione (68mg, 0.68mmol) and triethylamine (75mg, 0.75mmol) in dichloromethane (25ml) was stirred at room temperature for 18 hours. After this time the solvent was removed *in vacuo* and the residue was taken up in ethyl acetate (70ml). This solution was washed with 0.3M KHSO<sub>4</sub>, sat. NaHCO<sub>3</sub>, water and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by flash chromatography (eluant EtOAc:Pet. Ether 60-80°C; 7:3) yielding a colourless oil identified as the title compound (85mg, 0.29mmol, 47%).

MS:- ESI {M+H}<sup>+</sup> = 292.3

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 0.87-0.98 (6H, m), 1.19-1.25 (1H, m), 1.61-1.69 (2H, m), 1.84 (3H, s), 1.98 (3H, s), 2.15-2.25 (4H, m), 3.49-3.54 (1H, m), 3.62-3.69 (1H, m), 3.95-3.98 (1H, m), 4.75-4.79 (1H, m), 4.98 (1H, s), 11.09 (1H, d, J = 8.1Hz) ppm

**Example 5 – (2S)-1-(Glycylisoleucyl)pyrrolidine-2-carbonitrile**




**(a) (2S)-1-(Isoleucyl)pyrrolidine-2-carbonitrile**

To a solution of Boc-isoleucine hemihydrate (0.96g, 4mmol) and PyBOP® (2.34g, 4.5mmol) in dichloromethane (25ml) was added DIPEA (1.74ml, 10mmol). To that solution was added solid (S)-pyrrolidine-2-carbonitrile hydrochloride (0.60g, 4.5mmol) followed by another portion of DIPEA (697µl, 4mmol). The reaction mixture was stirred for 2h. The solvent was removed by rotary evaporation, and the residue was taken up in ethyl acetate. The resulting solution was washed with 0.3M sodium bisulfate (2x), saturated sodium bicarbonate (2x), water and saturated sodium chloride. The organic phase was dried with anhydrous sodium sulfate and the solvent was removed by rotary evaporation. The residue was dissolved in a mixture of TFA (95%) and water (5%). After 1 h most of TFA and water was removed under reduced pressure, and the residue was triturated with ether, resulting in the formation of a precipitate. The precipitate was collected and dried *in vacuo* to give the trifluoroacetate salt of the title product as a white solid; yield 0.58g (1.8mmol, 45%).

**(b) (2S)-1-(Glycylisoleucyl)pyrrolidine-2-carbonitrile**


To a solution of Boc-glycine (0.21g, 1.2mmol) and PyBOP® (0.62g, 1.2mmol) in dichloromethane (3ml) was added DIPEA (522µl, 3mmol). To that solution was added the product of Example 5a (0.28g, 0.9mmol) followed by another portion of DIPEA (157µl, 0.9mmol). The reaction mixture was stirred overnight. The solvent was removed by rotary evaporation and the residue was taken up in ethyl acetate. The resulting solution was washed with 0.3M sodium bisulfate (2x), saturated sodium bicarbonate (2x), water and saturated sodium chloride. The organic phase was dried with anhydrous sodium sulfate and the solvent was removed by rotary evaporation. The residue was dissolved in a mixture of TFA (95%) and water (5%) and the mixture was stirred overnight. Most of TFA and water was removed under reduced pressure. The residue was subjected to purification by reverse-phase HPLC to give the trifluoroacetate salt of the final product as a white powder; yield 171 mg (50%).



**Example 6 – (2S)-1-(Arginylisoleucyl)pyrrolidine-2-carbonitrile**

To a solution of Boc-Arg(Mtr)-OH (0.58g, 1.2mmol) and PyBOP® (0.62g, 1.2mmol) in dichloromethane (3ml) was added DIPEA (522μl, 3mmol). To that solution was added the product of Example 5a (0.28g, 0.9mmol) followed by another portion of DIPEA (157μl, 0.9mmol). The reaction mixture was stirred overnight. The solvent was removed by rotary evaporation, and the residue was taken up in ethyl acetate.

The resulting solution was washed with 0.3M sodium bisulfate (2x), saturated sodium bicarbonate (2x), water and saturated sodium chloride. The organic phase was dried with anhydrous sodium sulfate and the solvent was removed by rotary evaporation. The residue was dissolved in a mixture of TFA (95%) and water (5%) and the mixture was stirred overnight. Most of TFA and water was removed under reduced pressure and the residue was triturated with ether. The ethereal layer was decanted off, and the residue was subjected to purification by reverse-phase HPLC to give the trifluoroacetate salt of the final product as a white powder; yield 83mg (19%).


**Example 7 – (2S)-1-((2'S)-2'-(Acetoxymethoxycarbonylamino)-3',3'-dimethylbutanoyl)pyrrolidine-2-carbonitrile**

A solution of (2S)-1-((2'S)-2'-amino-3',3'-dimethylbutanoyl)pyrrolidine-2-carbonitrile hydrochloride (150mg, 0.61mmol; prepared according to Jenkins *et al.*, WO95/15309), acetoxymethyl *p*-nitrophenyl carbonate (168mg, 0.66mmol; prepared according to Alexander *et al.*, J. Med. Chem. 31, 318, 1988) and triethylamine (70mg, 0.70mmol) in dichloromethane (25ml) was stirred at room temperature for 18 hours. After this time the solvent was removed *in vacuo* and the residue was taken up in ethyl acetate (70ml). This solution was washed with 0.3M KHSO<sub>4</sub>, sat. NaHCO<sub>3</sub>, water and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by flash chromatography (eluant EtOAc:Pet. Ether 60-80°C; 4:6) yielding a white solid identified as the title compound (188mg, 0.58mmol, 95%).

MS:- ESI {M+H}<sup>+</sup> = 326.1

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.03 (9H, s), 2.09 (3H, s), 2.16-2.24 (4H, m), 3.72-3.77 (2H, m), 4.25 (1H, d, J = 9.6Hz), 4.77-4.80 (1H, m), 5.68 (1H, d), 5.68 (2H, s) ppm

**Example 8 – (2S)-1-((2'S)-2'-(1''-Acetoxyethoxycarbonylamino)-2'-cyclohexylacetyl)pyrrolidine-2-carbonitrile**




A solution of (2S)-1-((2'S)-2'-amino-2'-cyclohexylacetyl)pyrrolidine-2-carbonitrile trifluoroacetate (100mg, 0.28mmol; prepared according to Jenkins *et al.*, WO95/15309), α-acetoxyethyl *p*-nitrophenyl carbonate (76mg, 0.29mmol; prepared according to Alexander *et al.*, J. Med. Chem. 31, 318, 1988) and triethylamine (35mg, 0.35mmol) in dichloromethane (25ml) was stirred at room temperature for 18 hours. After this time the solvent was removed *in vacuo* and the residue was taken up in ethyl acetate (70ml). This solution was washed with 0.3M KHSO<sub>4</sub>, sat.

NaHCO<sub>3</sub>, water and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by flash chromatography (eluant EtOAc:Pet. Ether 60-80°C; 4:6) yielding a white solid identified as the title compound (43mg, 0.12mmol, 41%).

MS:- ESI {M+H}<sup>+</sup> = 366.2

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 0.97-1.21 (4H, m), 1.40-1.48 (3H, m), 1.67-1.77 (7H, m), 2.02 (3H, d, J=7.8Hz), 2.11-2.26 (4H, m), 3.65-3.73 (2H, m), 4.16-4.22 (1H, m), 4.76 (1H, d, J=4.2Hz), 5.36-5.41 (1H, m), 6.73-6.77 (1H, m) ppm

**Example 9 – (2S)-1-((2'S)-2'-(1''-Acetoxyethoxycarbonylamino)-4',4'-dimethylpentanoyl)pyrrolidine-2-carbonitrile**




A solution of (2S)-1-((2'S)-2'-amino-4',4'-dimethylpentanoyl)pyrrolidine-2-carbonitrile trifluoroacetate (100mg, 0.30mmol; prepared according to Jenkins *et al.*, WO95/15309), α-acetoxyethyl *p*-nitrophenyl carbonate (87mg, 0.33mmol; prepared according to Alexander *et al.*, *J. Med. Chem.* 31, 318, 1988) and triethylamine (40mg, 0.40mmol) in dichloromethane (25ml) was stirred at room temperature for 18 hours. After this time the solvent was removed *in vacuo* and the residue was taken up in ethyl acetate (70ml). This solution was washed with 0.3M KHSO<sub>4</sub>, sat. NaHCO<sub>3</sub>, water and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by flash chromatography (eluant EtOAc:Pet. Ether 60-80°C; 4:6) yielding a white solid identified as the title compound (32mg, 0.09mmol, 31%).

MS:- ESI {M+H}<sup>+</sup> = 354.2

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 0.97, 0.98 (9H, 2xs), 1.41-1.43 (3H, m), 1.44-1.62 (2H, m), 2.03 (3H, d, J=2.3Hz), 2.16-2.21 (4H, m), 3.61-3.63 (1H, m), 3.74-3.78 (1H, m), 4.45-4.52 (1H, m), 4.75-4.77 (1H, m), 5.24-5.29 (1H, m), 6.73-6.78 (1H, m) ppm

**Example 10 – (2S)-1-(N'-(1''-Acetoxyethoxycarbonyl)-O'-*tert*-butylserinyl)-pyrrolidine-2-carbonitrile**




A solution of (2S)-1-(O'-*tert*-butylserinyl)pyrrolidine-2-carbonitrile hydrochloride (30mg, 0.11mmol; prepared according to Jenkins *et al.*, WO95/15309),  $\alpha$ -acetoxyethyl *p*-nitrophenyl carbonate (32mg, 0.12mmol; prepared according to Alexander *et al.*, J. Med. Chem. 31, 318, 1988) and triethylamine (20mg, 0.20mmol) in dichloromethane (25ml) was stirred at room temperature for 18 hours. After this time the solvent was removed *in vacuo* and the residue was taken up in ethyl acetate (70ml). This solution was washed with 0.3M KHSO<sub>4</sub>, sat. NaHCO<sub>3</sub>, water and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by flash chromatography (eluant EtOAc:Pet. Ether 60-80°C; 4:6) yielding a white solid identified as the title compound (14mg, 0.038mmol, 35%).

MS:- ESI {M+H}<sup>+</sup> = 370.1

<sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  1.11-1.15 (9H, m), 1.41-1.45 (3H, m), 2.04 (3H, d, J = 4.9Hz), 2.10-2.15 (2H, m), 3.43-3.62 (5H, m), 3.90-4.00 (1H, m), 4.50-4.65 (1H, m), 4.73 (1H, d, J = 5.2Hz), 5.45-5.72 (1H, m), 6.76-6.79 (1H, m) ppm

Example 11 – (2S)-1-(N<sup>ω</sup>-(1'-Acetoxyethoxycarbonyl)-N<sup>ω</sup>-p-toluenesulphonyllysiny)pyrrolidine-2-carbonitrile




A solution of (2S)-1-(N<sup>ω</sup>-p-toluenesulphonyllysiny)pyrrolidine-2-carbonitrile trifluoroacetate (100mg, 0.20mmol; prepared according to Jenkins *et al.*, WO95/15309), α-acetoxyethyl p-nitrophenyl carbonate (61mg, 0.23mmol; prepared according to Alexander *et al.*, J. Med. Chem. 31, 318, 1988) and triethylamine (30mg, 0.30mmol) in dichloromethane (25ml) was stirred at room temperature for 18 hours. After this time the solvent was removed *in vacuo* and the residue was taken up in ethyl acetate (70ml). This solution was washed with 0.3M KHSO<sub>4</sub>, sat. NaHCO<sub>3</sub>, water and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by flash chromatography (eluant EtOAc:Pet. Ether 60-80°C; 7:3) yielding a white solid identified as the title compound (51mg, 0.10mmol, 49%).

MS:- ESI {M+H}<sup>+</sup> = 509.0

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.41-1.48 (6H, m), 1.51-1.69 (2H, m), 2.05 (3H, d, J = 18.3Hz), 2.12-2.28 (5H, m), 2.41 (3H, s), 2.86-2.93 (2H, m), 3.63-3.64 (2H, m), 4.38-4.42 (1H, m), 4.72-4.73 (1H, m), 4.74-4.79, 5.10-5.20 (1H, 2xm), 5.54-5.62 (1H, m), 6.74-6.79 (1H, m), 7.29 (2H, d, J = 7.7Hz), 7.71 (2H, d, J = 8.4Hz) ppm


**Example 12 – (2S)-1-(N-(1'-Acetoxyethoxycarbonyl)-N-(2''-(5'''-cyanopyridin-2'''-ylamino)ethyl)glyciny)pyrrolidine-2-carbonitrile**



A solution of 1-([2-[(5-cyanopyridin-2-yl)amino]ethyl]amino]acetyl)-2-cyano-(S)-pyrrolidine bis(trifluoroacetate) (100mg, 0.19mmol; prepared according to Villhauer *et al.*, WO98/19998),  $\alpha$ -acetoxyethyl *p*-nitrophenyl carbonate (56mg, 0.21mmol; prepared according to Alexander *et al.*, *J. Med. Chem.* **31**, 318, 1988) and triethylamine (50mg, 0.50mmol) in dichloromethane (25ml) was stirred at room temperature for 18 hours. After this time the solvent was removed *in vacuo* and the residue was taken up in ethyl acetate (70ml). This solution was washed with 0.3M KHSO<sub>4</sub>, sat. NaHCO<sub>3</sub>, water and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by flash chromatography (eluant EtOAc:Pet. Ether 60-80°C; 9:1) yielding a white solid identified as the title compound (13mg, 0.03mmol, 16%).  
MS:- ESI {M+H}<sup>+</sup> = 429.2

<sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  1.21-1.32 (3H, m), 1.40-1.46 (1H, m), 1.99-2.05 (4H, m), 2.17-2.31 (4H, m), 3.50-3.63 (6H, m), 4.40-4.50 (1H, m), 4.77 (1H, d, J = 5.9Hz), 6.45-6.49 (1H, m), 6.68-6.77 (1H, m), 7.44-7.48 (1H, m), 8.32 (1H, s) ppm

Example 13 - (2S)-1-(N'-(Benzyloxycarbonyl)-O'-*tert*-butylthreoninyl)-pyrrolidine-2-carbonitrile




A solution of (2S)-1-(O'-*tert*-butylthreoninyl)pyrrolidine-2-carbonitrile hydrochloride (35mg, 0.12mmol; prepared according to Jenkins *et al.*, WO95/15309), benzyl chloroformate (32mg, 0.13mmol) and triethylamine (24mg, 0.24mmol) in dichloromethane (25ml) was stirred at room temperature for 18 hours. After this time the solvent was removed *in vacuo* and the residue was taken up in ethyl acetate (70ml). This solution was washed with 0.3M KHSO<sub>4</sub>, sat. NaHCO<sub>3</sub>, water and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by flash chromatography (eluant: chloroform: methanol; 98:2) yielding a white solid identified as the title compound (47mg, 0.12mmol, 100%).

MS:- ESI {M+H}<sup>+</sup> = 388.3

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.10-1.30 (3H, m), 1.18 (9H, s), 2.00-2.45 (4H, m), 3.55-3.70 (1H, m), 3.85-4.00 (2H, m), 4.30-4.40 (1H, m), 4.70-4.80 (1H, m), 5.07 (2H, s), 5.75 (1H, d, J = 8.15Hz), 7.20-7.45 (5H, m) ppm

**Example 14 – (2S)-1-(S'-*tert*-Butyl-N'-(ethyloxycarbonyl)cysteiny)pyrrolidine-2-carbonitrile**




A solution of (2S)-1-(S'-*tert*-butylcysteiny)pyrrolidine-2-carbonitrile trifluoroacetate (1000mg, 0.27mmol; prepared according to Jenkins *et al.*, WO95/15309), ethylchloroformate (35mg, 0.32mmol) and triethylamine (50mg, 0.50mmol) in dichloromethane (25ml) was stirred at room temperature for 18 hours. After this time the solvent was removed *in vacuo* and the residue was taken up in ethyl acetate (70ml). This solution was washed with 0.3M KHSO<sub>4</sub>, sat. NaHCO<sub>3</sub>, water and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by flash chromatography (eluant EtOAc:Pet. Ether 60-80°C; 8:2) yielding a white solid identified as the title compound (30mg, 0.092mmol, 35%).

MS:- ESI {M+H}<sup>+</sup> = 328.1

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.18 (3H, t, J = 7 Hz), 1.30 (9H, s), 2.17-2.24 (4H, br m), 2.82-2.85 (2H, m), 3.70-3.82 (2H, br m), 4.05-4.09 (2H, m), 4.48-4.53 (1H, m), 4.74-4.77 (1H, m), 5.41-5.44 (1H, m) ppm.




**Example 15 – (2S)-1-(N<sup>ω</sup>-Acetyl-N<sup>α</sup>-benzoyllysinyl)pyrrolidine-2-carbonitrile**

A solution of (2S)-1-(N<sup>ω</sup>-acetyllysinyl)pyrrolidine-2-carbonitrile trifluoroacetate (100mg, 0.22mmol; prepared according to Jenkins *et al.*, WO95/15309), benzoyl chloride (343mg, 0.24mmol) and triethylamine (45mg, 0.45mmol) in dichloromethane (25ml) was stirred at room temperature for 18 hours. After this time the solvent was removed *in vacuo* and the residue was taken up in ethyl acetate (70ml). This solution was washed with 0.3M KHSO<sub>4</sub>, sat. NaHCO<sub>3</sub>, water and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by flash chromatography (eluant chloroform: methanol; 97:3) yielding a white solid identified as the title compound (83mg, 0.22mmol, 100%).

MS:- ESI {M+H}<sup>+</sup> = 387.6

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.56-1.78 (4H, br m), 1.94 (3H, s), 2.12-2.20 (4H, br m), 3.21-3.23 (2H, m), 3.59-3.72 (2H, m), 4.65-4.69 (2H, m), 5.07 (2H, s), 5.18-5.21 (1H, m), 6.69-6.72 (1H, m), 7.24-7.34 (5H, m) ppm

**Example 16 – (2S)-1-(N<sup>α</sup>-(Acetyl)-N<sup>ω</sup>-(benzyloxycarbonyl)ornithinyl)-pyrrolidine-2-carbonitrile**



A solution of (2S)-1-(N<sup>ω</sup>-(benzyloxycarbonyl)ornithinyl)pyrrolidine-2-carbonitrile trifluoroacetate (100mg, 0.23mmol; prepared according to Jenkins *et al.*, WO95/15309), acetyl chloride (20mg, 0.26mmol) and triethylamine (50mg, 0.50mmol) in dichloromethane (25ml) was stirred at room temperature for 18 hours. After this time the solvent was removed *in vacuo* and the residue was taken up in ethyl acetate (70ml). This solution was washed with 0.3M KHSO<sub>4</sub>, sat. NaHCO<sub>3</sub>, water and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by flash chromatography (eluant chloroform: methanol; 97:3) yielding a white solid identified as the title compound (49mg, 0.13mmol, 55%).

MS:- ESI {M+H}<sup>+</sup> = 371.2

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.30-1.65 (4H, m), 1.75-1.95 (2H, m), 1.90 (3H, s), 2.10-2.40 (4H, m), 3.10-3.30 (2H, m), 3.65-3.90 (2H, m), 4.70-4.90 (2H, m), 5.90-6.00 (1H, m), 7.30-7.50 (4H, m), 7.70-7.80 (2H, m) ppm

**Example 17 – *In vitro* inhibitory activity vs. DP-IV**

The compounds of the previous examples were assayed as inhibitors of DP-IV according to the method described in Ashworth *et al.* (Bioorg. Med. Chem. Lett. 1996, 6(10), 1163-66). No significant inhibitory activity was detected up to 10 $\mu$ M, indicating that the prodrugs of the invention are at least 1000 times less potent than the active inhibitors from which they are derived. Hence it can be assumed that any *in vivo* activity seen is due to bioconversion into the parent inhibitors.

**Example 18 – *In vivo* activity in glucose tolerance model**

The activity of the compounds was investigated in male Zucker Fatty Rats between 10 and 20 weeks of age. The animals were fasted overnight and then administered with the test compound (10mg/kg) as a solution by oral gavage. One hour later a blood sample (200 $\mu$ l) is taken from the tail vein to establish a baseline (t = 0) glucose level, then the animals are given glucose (1g/kg as a 40% wt/vol solution) orally. Further blood samples are taken at t = 10, 20, 30, 60 and 120 minutes. Glucose is determined by an enzymatic assay. Typical results are given in the Table below.

Compound	Blood glucose (mg/dl); mean $\pm$ SE, n = 4					
	t = 0	t = 10	t = 20	t = 30	t = 60	t = 120
vehicle	95.1 $\pm$ 7.36	151.6 $\pm$ 8.12	164.3 $\pm$ 10.7	153.2 $\pm$ 7.8	153.4 $\pm$ 7.8	122.4 $\pm$ 7.0
Example 2	80.2 $\pm$ 4.2	122.2 $\pm$ 7.8	117.8 $\pm$ 4.8	104.4 $\pm$ 5.6	117.6 $\pm$ 6.7	111.3 $\pm$ 12.1
Example 3	86.4 $\pm$ 3.4	175.1 $\pm$ 3.8	148.5 $\pm$ 23.2	136.7 $\pm$ 16.9	120.6 $\pm$ 8.3	101.9 $\pm$ 4.5
Example 5	80.5 $\pm$ 1.4	141.3 $\pm$ 14.5	134.2 $\pm$ 10.3	129.2 $\pm$ 8.2	114.6 $\pm$ 8.1	121.2 $\pm$ 5.8
Reference	91.4 $\pm$ 5.2	125.8 $\pm$ 8.9	110.0 $\pm$ 21.5	110.9 $\pm$ 4.4	112.2 $\pm$ 7.5	108.7 $\pm$ 8.9

The reference compound in the above experiments was the compound of Example 11 of WO95/15309. This is the parent compound from which the prodrugs of Examples 2 – 6 of the present disclosure are derived.

It is clear from the above results that the prodrugs are effective at reducing hyperglycaemia after the glucose challenge, but that they are not always as effective as the reference compound at early time points. This is what would be expected for prodrugs that are converted in high yield to the parent drug. The results at early time points are a result of the need for metabolic conversion of the circulating prodrug.

In a separate experiment the test compound was given at the same dose (10mg/kg) but 12 hours before the oral glucose challenge. The results are given below.

Compound	Blood glucose (mg/dl); mean $\pm$ SE, n = 4						
	t = 0	t = 10	t = 20	t = 30	t = 60	t = 120	AUC
vehicle	84.2 $\pm$ 3.7	145.5 $\pm$ 6.6	134.3 $\pm$ 8.0	127.2 $\pm$ 10.1	122.9 $\pm$ 8.7	112.2 $\pm$ 8.6	4556 $\pm$ 458
Example 1	83.7 $\pm$ 3.8	113.5 $\pm$ 10.8	111.3 $\pm$ 9.9	91.9 $\pm$ 11.8	99.3 $\pm$ 10.0	116.9 $\pm$ 14.4	2430 $\pm$ 591

The AUC (area under the concentration-time curve) is greatly reduced, showing that the prodrug enables significant antihyperglycaemic activity to be maintained for 12 hours.

The above results illustrate that the compounds of the present invention exhibit antihyperglycaemic activity after oral administration in a relevant animal model of glucose intolerance. Hence it is to be expected that they would be effective in the treatment of human impaired glucose tolerance and type 2 diabetes. Furthermore, the in vivo results confirm that the prodrugs are converted to active DP-IV inhibitors in the circulation, and that they could be used in the treatment of all the other pathologies for which such inhibitors have been proposed as therapeutic agents.

**Example 19 - Pharmaceutical Formulations**19A - 50mg Tablets

Tablets containing 50mg of the compound of Example 1 are prepared from the following:

Compound of Example	154.5g
Corn Starch	53.5g
Hydroxypropylcellulose	13.5g
Carboxymethylcellulose calcium	11.0g
Magnesium stearate	2.0g
Lactose	165.5g
<hr/>	
Total	400.0g

The materials are blended and then pressed to give 2000 tablets of 200mg, each containing 50mg of the compound of Example 1.

The compounds of Examples 2, 3 and 5 were formulated separately into respective tablets in the same manner. The compounds of Examples 4 and 6 to 16 were similarly formulated separately into tablets containing 100mg of the respective compounds.

19B - 100mg Suppository


Suppositories containing 100mg of the compound of Example 2 are prepared from the following:

Compound of Example 2	154.5g
Corn Starch	210.0g
Colloidal silica	2.5g
Povidone 30	49.0g
Magnesium stearate	23.0g
Adipic acid	57.0g
Sodium bicarbonate	43.0g
Sodium lauryl sulphate	5.0g
Lactose	456.0g
<hr/>	
Total	1000.0g


The materials are blended and then pressed to give suppositories of 1g. each containing 100mg of compound of Example 2. The compounds of Examples 1, 3 to 5 and 6 to 16 were formulated into respective suppositories in the same manner.

## CLAIMS

1. A compound according to the general formula 1, or a pharmaceutically acceptable salt thereof,



wherein:



$R^1$  is selected from H,  $C_1 - C_8$  alkyl (including branched alkyl and cycloalkyl),  $(CH_2)_aNHW^1$ ,  $(CH_2)_bCOW^2$ ,  $(CH_2)_cOW^3$ ,  $CH(Me)OW^4$ ,  $(CH_2)_d-C_6H_4-W^5$  and  $(CH_2)_eSW^6$ , where a is 2-5, b is 1-4, c is 1-2, d is 1-2, e is 1-3,  $W^1$  is  $COW^6$ ,  $CO_2W^6$  or  $SO_2W^6$ ,  $W^2$  is OH,  $NH_2$ ,  $OW^6$  or  $NHW^6$ ,  $W^3$  is H or  $W^6$ ,  $W^4$  is H or  $W^6$ ,  $W^5$  is H, OH or OMe, and  $W^6$  is  $C_1 - C_8$  alkyl, benzyl, optionally substituted phenyl or optionally substituted heteroaryl where the optional substituents are up to two groups selected from  $C_1 - C_3$  alkyl,  $C_1 - C_3$  alkoxy, F and Cl;

$R^2$  is H or  $-(CH_2)_nNH-C_6H_4N-Y$ , where n is 2-4 and Y is selected from H, F, Cl,  $NO_2$  and CN, or

$R^1$  and  $R^2$  together are  $-(CH_2)_p-$ , where p is 3 or 4;

X is selected from

- (i) L-alpha-amino acyl groups selected from Ala, Arg, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Tyr and Val,
- (ii) groups  $R^3CO$ , where  $R^3$  is H,  $C_1 - C_6$  alkyl (including branched alkyl and cycloalkyl), or phenyl,
- (iii) groups  $R^4COOC(R^5)(R^6)OCO$ , where  $R^4$  is H,  $C_1 - C_6$  alkyl (including branched alkyl and cycloalkyl), benzyl or optionally substituted phenyl where the substituents are up to two groups selected from  $C_1 - C_3$  alkyl,  $C_1 - C_3$  alkoxy, F and Cl, and  $R^5$  and  $R^6$  are independently H or  $C_1 - C_6$  alkyl, or  $R^5$  and  $R^6$  together are  $-(CH_2)_m-$  where m is an integer of 4-6, and
- (iv) methoxycarbonyl, ethoxycarbonyl and benzyloxycarbonyl groups;

$R^7$  is selected from pyridyl and optionally substituted phenyl, where the substituents are up to two groups selected from  $C_1 - C_3$  alkyl,  $C_1 - C_3$  alkoxy, F, Cl,  $NO_2$ , CN and  $CO_2H$ ;

$R^8$  is H or  $C_1 - C_3$  alkyl; and

$R^9$  is H,  $C_1 - C_6$  alkyl, phenyl or  $C_1 - C_6$  alkoxy.

2. A compound according to Claim 1 wherein  $R^1$  is other than H and  $R^2$  is H.
3. A compound according to Claim 2 wherein  $R^1$  is  $C_1 - C_6$  alkyl.
4. A compound according to Claim 1 wherein  $R^1$  is H, A is a group according to general formulae 2 or 4, and  $R^2$  is  $-(CH_2)_nNH-C_5H_3N-Y$ .
5. A compound according to Claim 4 wherein n is 2 and Y is CN.



6. A compound according to Claim 5 wherein the NH substituent is at the 2-position and the cyano group is at the 5-position of the pyridyl ring.
7. A compound according to Claim 1 wherein A is a group according to general formula 2 and X is an amino acyl group.
8. A compound according to Claim 7 wherein the amino acyl group is a lysyl or argininy group.
9. A compound according to Claim 8 wherein the amino acyl group is an argininy group.
10. A compound according to Claim 7 wherein the amino acyl group is a glyciny group.
11. A compound according to Claim 1 wherein A is a group according to general formula 2 and X is  $R^4COOC(R^5)(R^6)OCO$ .
12. A compound according to Claim 11 wherein  $R^4$  is  $C_1 - C_6$  alkyl.
13. A compound according to Claim 11 wherein one of  $R^5$  and  $R^6$  is H and the other is methyl.
14. A compound according to Claim 11 wherein  $R^4$  is methyl, one of  $R^5$  and  $R^6$  is H and the other is methyl.
15. A compound according to Claim 1 wherein A is a group according to general formula 2 and X is methoxycarbonyl.
16. A compound according to Claim 1 wherein A is a group according to general formula 3.

17. A compound according to Claim 1 wherein A is a group according to general formula 4.
18. A compound according to Claim 17 wherein R<sup>8</sup> is C<sub>1</sub> – C<sub>3</sub> alkyl.
19. A compound according to Claim 18 wherein R<sup>8</sup> is methyl.
20. A compound according to Claim 17 wherein R<sup>9</sup> is C<sub>1</sub> – C<sub>3</sub> alkyl or C<sub>1</sub> – C<sub>3</sub> alkoxy.
21. A compound according to Claim 20 wherein R<sup>9</sup> is methyl or methoxy.
22. A compound according to Claim 1 selected from
  - (2S)-1-((2'S)-2'-(1''-acetoxyethoxycarbonylamino)-3',3'-dimethylbutanoyl)-pyrrolidine-2-carbonitrile;
  - (2S)-1-(N'-(1''-acetoxyethoxycarbonyl)isoleucyl)pyrrolidine-2-carbonitrile;
  - (2S)-1-(N'-(methoxycarbonyl)isoleucyl)pyrrolidine-2-carbonitrile;
  - (2S)-1-((N')-(4''-oxopent-2''-en-2''-yl)isoleucyl)pyrrolidine-2-carbonitrile;
  - (2S)-1-(glycylisoleucyl)pyrrolidine-2-carbonitrile;
  - (2S)-1-(arginylisoleucyl)pyrrolidine-2-carbonitrile;
  - (2S)-1-((2'S)-2'-(acetoxymethoxycarbonylamino)-3',3'-dimethylbutanoyl)-pyrrolidine-2-carbonitrile;
  - (2S)-1-((2'S)-2'-(1''-acetoxyethoxycarbonylamino)-2'-cyclohexylacetyl)-pyrrolidine-2-carbonitrile;
  - (2S)-1-((2'S)-2'-(1''-acetoxyethoxycarbonylamino)-4',4'-dimethylpentanoyl)-pyrrolidine-2-carbonitrile;
  - (2S)-1-(N'-(1''-acetoxyethoxycarbonyl)-O'-*tert*-butylserinyl)pyrrolidine-2-carbonitrile;
  - (2S)-1-(N<sup>α</sup>-(1'-acetoxyethoxycarbonyl)-N<sup>ω</sup>-*p*-toluenesulphonyllysiny)l)-pyrrolidine-2-carbonitrile;

(2S)-1-(N-(1'-acetoxyethoxycarbonyl)-N-(2''-(5'''-cyanopyridin-2'''-ylamino)-ethyl)glyciny)pyrrolidine-2-carbonitrile;  
(2S)-1-(N'-(benzyloxycarbonyl)-O'\_*tert*-butylthreoniny)pyrrolidine-2-carbonitrile;  
(2S)-1-(S'-*tert*-butyl-N'-(ethyloxycarbonyl)cysteiny)pyrrolidine-2-carbonitrile;  
(2S)-1-(N<sup>ω</sup>-acetyl-N<sup>α</sup>-benzoyllysiny)pyrrolidine-2-carbonitrile; and  
(2S)-1-(N<sup>α</sup>-(acetyl)-N<sup>ω</sup>-(benzyloxycarbonyl)ornithiny)pyrrolidine-2-carbonitrile.

23. At least one compound selected from optical isomers of compounds according to any preceding claim.
24. At least one compound selected from pharmaceutically acceptable salts of compounds according to any preceding claim.
25. A pharmaceutical composition containing agent selected from compounds according to any preceding Claim.
26. A composition according to Claim 25, which composition is for the treatment of impaired glucose tolerance or type 2 diabetes.
27. A method of treatment of impaired glucose tolerance or type 2 diabetes which comprises the administration to a person in need of such treatment of a therapeutically effective amount of compound(s) according to any of Claims 1-24.
28. The use, in the preparation of a medicament for the treatment of impaired glucose tolerance or type 2 diabetes, of a compound or compounds according to any of Claims 1 to 24.

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
7 June 2001 (07.06.2001)

PCT

(10) International Publication Number  
**WO 01/40180 A3**

(51) International Patent Classification<sup>7</sup>: C07D 207/16,  
A61K 31/401

(21) International Application Number: PCT/GB00/04572

(22) International Filing Date:  
30 November 2000 (30.11.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
9928330.1 30 November 1999 (30.11.1999) GB

(71) Applicant (for all designated States except US): FER-  
RING BV [NL/NL]; Polaris Avenue 144, NL-2132 JX  
Hoofddorp (NL).

(72) Inventor; and

(75) Inventor/Applicant (for US only): EVANS, David,  
Michael [GB/GB]; 114 Adelaide Road, St. Denys,  
Southampton SO17 2HX (GB).

(74) Agent: GEERING, Keith, Edwin; Reddie & Grose, 16  
Theobalds Road, London WC1X 8PL (GB).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).


Published:

— with international search report

(88) Date of publication of the international search report:  
8 November 2001

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: N-SUBSTITUTED 2-CYANOPYRROLIDINES AND THEIR USE AS ANTIDIABETIC AGENTS



(57) Abstract: Compounds which are 1-(2'-aminoacyl)-2-cyanopyrrolidine derivatives according to general formula (1) are DP-IV inhibitors for treatment of impaired glucose tolerance or type 2 diabetes; wherein A is selected from groups (2, 3 and 4); X is selected from aminoacyl groups corresponding to the natural amino acids, acyl groups R<sup>3</sup>CO, groups R<sup>4</sup>COOC(R<sup>5</sup>)(R<sup>6</sup>)OCO, methoxycarbonyl, ethoxycarbonyl and benzyloxycarbonyl; R<sup>1</sup> is selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl residues, (CH<sub>2</sub>)<sub>a</sub>NHW<sup>1</sup>, (CH<sub>2</sub>)<sub>b</sub>COW<sup>2</sup>, (CH<sub>2</sub>)<sub>c</sub>OW<sup>3</sup>, CH(Me)OW<sup>4</sup>, (CH<sub>2</sub>)<sub>d</sub>-C<sub>6</sub>H<sub>4</sub>-W<sup>5</sup> and (CH<sub>2</sub>)<sub>e</sub>SW<sup>6</sup>, where a is 2-5, b is 1-4, c is 1-2, d is 1-2, e is 1-3, W<sup>1</sup> is COW<sup>6</sup>, CO<sub>2</sub>W<sup>6</sup> or SO<sub>2</sub>W<sup>6</sup>, W<sup>2</sup> is OH, NH<sub>2</sub>, OW<sup>6</sup> or NHW<sup>6</sup>, W<sup>3</sup> is H or W<sup>6</sup>, W<sup>4</sup> is H or W<sup>6</sup>, W<sup>5</sup> is H, OH or OMe, and W<sup>6</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, optionally substituted phenyl, optionally substituted heteroaryl or benzyl and R<sup>2</sup> is selected from H and (CH<sub>2</sub>)<sub>n</sub>-C<sub>5</sub>H<sub>3</sub>N-Y, where n is 2-4 and Y is H, F, Cl, NO<sub>2</sub> or CN, or R<sup>1</sup> and R<sup>2</sup> together are -(CH<sub>2</sub>)<sub>p</sub>-where p is 3 or 4; R<sup>3</sup> is selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl and phenyl; R<sup>4</sup> is selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl and optionally substituted phenyl; R<sup>5</sup> and R<sup>6</sup> are each independently selected from H and C<sub>1</sub>-C<sub>6</sub> alkyl or together are -(CH<sub>2</sub>)<sub>m</sub>-, where m is 4-6; R<sup>7</sup> is selected from pyridyl and optionally substituted phenyl; R<sup>8</sup> is selected from H and C<sub>1</sub>-C<sub>3</sub> alkyl; and R<sup>9</sup> is selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy and phenyl.



WO 01/40180 A3

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/GB 00/04572

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D207/16 A61K31/401

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data, BEILSTEIN Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97 40832 A (PAULY ROBERT P ; PEDERSON RAY A (CA); MCINTOSH CHRISTOPHER H S (CA)) 6 November 1997 (1997-11-06) cited in the application the whole document ---	1-28
Y	WO 95 15309 A (FERRING BV ; JENKINS PAUL D (GB); JONES D MICHAEL (GB); SZELKE MICH) 8 June 1995 (1995-06-08) cited in the application see page 32 and example 1 the whole document ---	1-28
Y	WO 98 19998 A (CIBA GEIGY AG ; VILLHAUER EDWIN BERNARD (US)) 14 May 1998 (1998-05-14) cited in the application the whole document ---	1-28
-/-		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

23 February 2001

Date of mailing of the international search report

23. 05. 01

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Scruton-Evans, I

## INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/GB 00/04572

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	HUGHES, THOMAS E. ET AL: "NVP-DPP728 (1-[[[2-[(5-Cyanopyridin-2-yl)amino]ethyl]amino]acetyl]-2-cyan (S)-pyrrolidine), a Slow-Binding Inhibitor of Dipeptidyl Peptidase IV" BIOCHEMISTRY (1999), 38(36), 11597-11603, XP000983860 the whole document ---	1-28
X	YAMADA, TAKASHI ET AL: "Studies of unusual amino acids and their peptides. VIII. The syntheses of an iminohexapeptide as a model of bottromycin and its related iminopeptides" BULL. CHEM. SOC. JPN. (1977), 50(7), 1827-30, XP000983920 see compounds VI and X ---	1-3,23
X	YAMADA, TAKASHI ET AL: "Studies of unusual amino acids and their peptides. IX. The synthetic study of bottromycins B1 and B2" BULL. CHEM. SOC. JPN. (1978), 51(3), 878-83, XP000983921 see compound RN66854-31-7, compound 6 ---	1,3,23
P,Y	WO 00 56297 A (BROQUA PIERRE ;FERRING BV (NL)) 28 September 2000 (2000-09-28) see whole document and example 1A ---	1-28
P,Y	WO 00 56296 A (BROQUA PIERRE ;FERRING BV (NL)) 28 September 2000 (2000-09-28) see whole document, especially example 1A ---	1-28
P,X	WO 00 55125 A (BRYANT CLIFFORD M ;BUNIN BARRY A (US); AXYS PHARM INC (US); KRAYNA) 21 September 2000 (2000-09-21) see formula I and definitions of R2 and R4 ---	1-28
E	US 6 172 081 B1 (R DAMON) 9 January 2001 (2001-01-09) the whole document ---	1-28
P,Y	WO 00 34241 A (NOVARTIS ERFIND VERWALT GMBH ;NOVARTIS AG (CH); VILLHAUER EDWIN BE) 15 June 2000 (2000-06-15) the whole document ---	1-28

-/--

INTERNATIONAL SEARCH REPORT

Intern: al Application No  
PCT/GB 00/04572

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>ASHWORTH D M ET AL: "2-Cyanopyrrolidides as potent, stable inhibitors of dipeptidyl peptidase IV" BIOORGANIC &amp; MEDICINAL CHEMISTRY LETTERS, OXFORD, GB, vol. 6, no. 10, 21 May 1996 (1996-05-21), pages 1163-1166, XP004134889 ISSN: 0960-894X the whole document -----</p>	1-28



# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/GB 00/04572

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claim 27 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



# INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. al Application No

PCT/GB 00/04572

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9740832 A	06-11-1997	DE 19616486 A AU 721477 B AU 3023397 A CA 2252576 A CN 1216468 A EP 0896538 A NZ 332707 A	30-10-1997 06-07-2000 19-11-1997 06-11-1997 12-05-1999 17-02-1999 28-10-1999
WO 9515309 A	08-06-1995	AU 1113395 A AU 8421998 A CA 2178066 A CN 1141033 A CZ 9601595 A EP 0731789 A FI 962315 A HU 76274 A JP 9509921 T NO 962269 A PL 314838 A US 5939560 A ZA 9409525 A	19-06-1995 12-11-1998 08-06-1995 22-01-1997 15-01-1997 18-09-1996 05-08-1996 28-07-1997 07-10-1997 30-07-1996 30-09-1996 17-08-1999 02-08-1995
WO 9819998 A	14-05-1998	AU 726186 B AU 5318498 A BR 9714130 A CN 1236361 A CZ 9901615 A EP 0937040 A HU 0000323 A JP 2000511559 T NO 992028 A PL 332777 A SK 60899 A TR 9901004 T	02-11-2000 29-05-1998 29-02-2000 24-11-1999 11-08-1999 25-08-1999 28-08-2000 05-09-2000 28-04-1999 11-10-1999 10-04-2000 21-07-1999
WO 0056297 A	28-09-2000	AU 3318600 A	09-10-2000
WO 0056296 A	28-09-2000	AU 3317700 A	09-10-2000
WO 0055125 A	21-09-2000	AU 3746100 A AU 3748600 A WO 0055126 A	04-10-2000 04-10-2000 21-09-2000
US 6172081 B		NONE	
WO 0034241 A	15-06-2000	AU 1658000 A US 6166063 A	26-06-2000 26-12-2000